

Amendment to the Specification

Please replace the paragraph beginning on page 20, lines 9-20 with the following paragraph:

Without being bound by a specific theory or mechanism, the following events likely occur. Upon inoculation, fibroblasts likely form a physiological attachment to the collagen matrix by binding via collagen-specific receptors. Because the matrix is reticulated and thus contains multiple continuous surfaces, as opposed to being perforated with direct channels or openings from a top surface to a bottom surface, the fibroblasts or other dermal cells being inoculated need not fill these channels or openings in the matrix before the epidermal cells may be added. Rather, upon inoculation, the dermal cells attach to the reticulations, and thus are able to provide a continuous surface lamination for the ~~subsequently~~ subsequent inoculation of epidermal cells within a shorter time period than is possible using a perforated matrix.

BEST AVAILABLE COPY

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.